NUMAL: Lidocaine Gel BP 2%

Prescription Required

Name PSP RESEARCH PVT I LTD
Address: GURUGRAM, HY INDIA
SALT COMPOSITION

Lidocaine (2% w/w) + Chlorhexidine Gluconate (1% w/w) + Metronidazole (1% w/w)

STORAGE
Store below 30°C

PRODUCT INTRODUCTION

NUMAL Forte Gel is a combination medicine used in the treatment of mouth ulcers. It promotes the healing of mouth ulcers by forming a protective coating over the mouth sores. It also blocks the transmission of pain signals to the brain to decrease the sensation of pain.

NUMAL Forte Gel should be used strictly as advised by your doctor. Use it in the dose and duration as directed by your doctor. You should not use more quantity or more often than prescribed by your doctor. This will not make it work any faster rather increase the chance of side effects.

Using of this medicine may cause some common side effects such as application site reactions (burning, irritation, itching, and redness), dryness in mouth, metallic taste, nausea, and staining of teeth. These are usually temporary and resolve with time. If they do not get better with time or worsen, you should let your doctor know. Your doctor may be able to suggest ways of preventing or reducing the side effects.

Before using the medicine, it is important to tell your doctor if you are using or have recently used any other medicines for the same condition or other diseases. To make sure the medicine is safe, pregnant and breastfeeding women should consult their doctor before using the medicine.

USES OF NUMAL DENTAL GEL

Treatment of Mouth ulcers

BENEFITS OF NUMAL DENTAL GEL

In Treatment of Mouth ulcers

Mouth ulcers are small painful sores on inner lips, gums, tongue, roof of the mouth or throat that may interfere with eating, drinking and even talking. NUMAL Forte Gel blocks chemicals in our brain that are responsible for pain sensation and inflammation. Therefore it provides relief from pain, discomfort and symptoms of inflammation such as redness or swelling due to mouth ulcers. NUMAL Forte Gel also contains antimicrobial agents that kill the harmful bacteria and prevent any further infections. This makes it easier for you to go about your daily activities. Continue using it as advised by the doctor to get maximum benefit.

SIDE EFFECTS OF NUMAL DENTAL GEL

Most side effects do not require any medical attention and disappear as your body adjusts to the medicine. Consult your doctor if they persist or if you're worried about them

Common side effects of NUMAL

- Application site reactions (burning, irritation, itching and redness)
- Dryness
- Metallic taste

- Nausea
- Staining of teeth

HOW TO USE NUMAL DENTAL GEL

This medicine is for external use only. Use this medicine in the dose and duration as advised by your doctor.

HOW NUMAL DENTAL GEL WORKS

NUMAL Forte Gel is a combination of three medicines: Lidocaine, Chlorhexidine Gluconate and Metronidazole, which treats mouth sores. Lidocaine is a local anesthetic which forms a protective coating over the mouth sores and blocks pain signals to further relieve pain. Chlorhexidine Gluconate is an antiseptic and disinfectant which effectively cleans out the ulcer surface to prevent infection. Metronidazole is an antibiotic which kills the bacteria and other microorganisms by damaging their DNA.

SAFETY ADVICE



Alcohol

No interaction found/established



Pregnancy

CONSULT YOUR DOCTOR

NUMAL Forte Gel may be unsafe to use during pregnancy. Although there are limited studies in humans, animal studies have shown harmful effects on the developing baby. Your doctor will weigh the benefits and any potential risks before prescribing it to you. Please consult your doctor.



Breast feeding

SAFE IF PRESCRIBED

NUMAL Forte Gel is probably safe to use during breastfeeding. Limited human data suggests that the drug does not represent any significant risk to the baby.



Driving

No interaction found/established



Kidney

No interaction found/established



Liver

No interaction found/established

WHAT IF YOU FORGET TO TAKE NUMAL DENTAL GEL?

If you miss a dose of NUMAL Forte Gel, use it as soon as possible. However, if it is almost time for your next dose, skip the missed dose and go back to your regular schedule. Do not double the dose.